## Foliations with singularities

A foliation with singularities *F* of a 2-manifold *M*, for our purposes, is a foliation of *M* where there are finitely many points x<sub>1</sub>,..., x<sub>l</sub> ∈ *M* at which some number p = p(x<sub>k</sub>) = p<sub>k</sub> ≥ 3 of the leaves meet (these are the prongs of the singularity):



Thermodynamics of Pseudo-Anosov Diffeomorphisms

Dominic Veconi

Pseudo-Anosov Maps

Smooth Ergodic Theory

Young Towers

# Pseudo-Anosov Homeomorphisms

- A homeomorphism f : M → M of a 2-manifold M is pseudo-Anosov if there are two f-invariant foliations with singularities, F<sup>s</sup> and F<sup>u</sup>, for which:
  - the foliations share the same singularities, and the same number of prongs;
  - the foliations intersect transversally away from the singularities;
  - 3. there is a  $\lambda > 1$  such that for x, y in the same  $\mathcal{F}^{s}$ -leaf,  $\rho^{s}(f(x), f(y)) = \lambda^{-1}\rho^{s}(x, y)$ , and for x, y in the same  $\mathcal{F}^{u}$ -leaf,  $\rho^{u}(f(x), f(y)) = \lambda \rho^{u}(x, y)$ ;

where  $\rho^s$  and  $\rho^u$  are the distances in the  $\mathcal{F}^s$  and  $\mathcal{F}^u$  foliations with respect to a Riemannian metric on M that has a density vanishing at the singularities.

Thermodynamics of Pseudo-Anosov Diffeomorphisms

Dominic Veconi

Pseudo-Anosov Maps

Smooth Ergodic Theory

Young Towers

# Pseudo-Anosov Homeomorphisms



Thermodynamics of Pseudo-Anosov Diffeomorphisms

Dominic Veconi

Pseudo-Anosov Maps

Smooth Ergodic Theory

Young Towers

Proof of Main Result

The blue curves represent the stable foliation, along which nearby points contract; and the red curves represent the unstable foliation, along which nearby points expand.

# Nielsen-Thurston Classification

#### Theorem (Nielsen, Thurston)

Any homeomorphism on a compact topological manifold M is isotopic to a map f that is one of the following:

- f is periodic: there is a positive integer m with f<sup>m</sup> = Id;
  - EG.  $f = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} : \mathbb{T}^2 \to \mathbb{T}^2;$
- f is reducible: there is a closed curve on M that is f-invariant (these are also known as Dehn twists);
  - EG.  $f = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} : \mathbb{T}^2 \to \mathbb{T}^2;$
- f is pseudo-Anosov;
  - EG.  $f = \begin{pmatrix} 2 & 1 \\ 1 & 1 \end{pmatrix} : \mathbb{T}^2 \to \mathbb{T}^2$ .

The pseudo-Anosov maps form an open set in the homeomorphisms of M, and exhibit the most interesting dynamical properties.

Thermodynamics of Pseudo-Anosov Diffeomorphisms

Dominic Veconi

Pseudo-Anosov Maps

Smooth Ergodic Theory

Young Towers

# Examples from Anosov maps

- An Anosov diffeomorphism is a pseudo-Anosov homeomorphism with no singularities.
- ► Linear Anosov maps on T<sup>2</sup> lift to pseudo-Anosov maps on higher-genus surfaces via branched coverings (may be necessary to lift powers of Anosov maps).



Thermodynamics of Pseudo-Anosov Diffeomorphisms

Dominic Veconi

Pseudo-Anosov Maps

Smooth Ergodic Theory

Young Towers

▶ Orientable genus-2 surface S<sub>2</sub> can be horizontally split into two cylinders C<sub>1</sub> and C<sub>2</sub>, each of which admits a Dehn twist T<sub>1</sub> and T<sub>2</sub> resp. Note dT<sub>1</sub> = (<sup>1</sup><sub>0</sub><sup>1</sup><sub>1</sub>) on C<sub>1</sub>, and dT<sub>2</sub> = (<sup>1</sup><sub>0</sub><sup>2</sup><sub>1</sub>) on C<sub>2</sub>, away from the identified vertex (singularity).



Thermodynamics of Pseudo-Anosov Diffeomorphisms

Dominic Veconi

Pseudo-Anosov Maps

Smooth Ergodic Theory

Young Towers

► The Dehn twists T<sub>1</sub> and T<sub>2</sub> can be composed to form a horizontal "multi-twist" T := T<sub>1</sub><sup>2</sup> ∘ T<sub>2</sub>, whose differential away from the vertex is (<sup>1</sup><sub>0</sub> <sup>2</sup><sub>1</sub>):



Thermodynamics of Pseudo-Anosov Diffeomorphisms

Dominic Veconi

Pseudo-Anosov Maps

Smooth Ergodic Theory

Young Towers

► A similar vertical multi-twist S can be defined on the cylinder made from the red and blue squares, and the cylinder made from the green square. The corresponding differential is (<sup>1</sup>/<sub>2</sub> <sup>0</sup>/<sub>1</sub>).



Thermodynamics of Pseudo-Anosov Diffeomorphisms

Dominic Veconi

Pseudo-Anosov Maps

Smooth Ergodic Theory

Young Towers

- $T \circ S$  has the constant differential  $\begin{pmatrix} 1 & 2 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 2 & 1 \end{pmatrix} = \begin{pmatrix} 5 & 2 \\ 2 & 1 \end{pmatrix}$ .
- ► Eigenvalues are  $3 2\sqrt{2}$  and  $3 + \sqrt{2}$ , with resp. eigenvectors  $\begin{pmatrix} 1-\sqrt{2} \\ 1 \end{pmatrix}$  and  $\begin{pmatrix} 1+\sqrt{2} \\ 1 \end{pmatrix}$ .
- ► This is a pseudo-Anosov map whose stable/unstable foliations are parallel to the eigendirections for 3 2√2 and 3 + 2√2, resp. The vertex is a 6-pronged singularity (unstable prongs illustrated:)



Thermodynamics of Pseudo-Anosov Diffeomorphisms

Dominic Veconi

Pseudo-Anosov Maps

Smooth Ergodic Theory

Young Towers

- $T \circ S$  has the constant differential  $\begin{pmatrix} 1 & 2 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 2 & 1 \end{pmatrix} = \begin{pmatrix} 5 & 2 \\ 2 & 1 \end{pmatrix}$ .
- Eigenvalues are  $3 2\sqrt{2}$  and  $3 + \sqrt{2}$ , with resp. eigenvectors  $\begin{pmatrix} 1-\sqrt{2} \\ 1 \end{pmatrix}$  and  $\begin{pmatrix} 1+\sqrt{2} \\ 1 \end{pmatrix}$ .
- ► This is a pseudo-Anosov map whose stable/unstable foliations are parallel to the eigendirections for 3 2√2 and 3 + 2√2, resp. The vertex is a 6-pronged singularity (unstable prongs illustrated:)



▶ In fact, this map is the lift of the linear Anosov map on  $\mathbb{T}^2$  induced by  $\begin{pmatrix} 5 & 2 \\ 2 & 1 \end{pmatrix}$ .

Thermodynamics of Pseudo-Anosov Diffeomorphisms

Dominic Veconi

Pseudo-Anosov Maps

Smooth Ergodic Theory

Young Towers

## Properties

Suppose  $f: M \to M$  is a pseudo-Anosov homeomorphism with expansion factor  $\lambda > 1$ .

- There is a Riemannian metric on *M* inducing a volume ν under which *f* is invariant. Furthermore, if *U* is a neighborhood of a singularity x<sub>i</sub> ∈ *M* and φ : U → ℝ<sup>2</sup> is a coordinate chart, ν has a density with respect to φ<sub>\*</sub><sup>-1</sup>(Leb) vanishing at x<sub>i</sub>.
- f admits a finite Markov partition, with respect to which f is Bernoulli.
- If x ∈ M is not a singularity, then f is smooth at x and there are orthonormal bases of T<sub>x</sub>M and T<sub>f(x)</sub>M with respect to which df<sub>x</sub> has the matrix form (<sup>λ</sup><sub>0</sub> 0/<sub>0</sub>).

Thermodynamics of Pseudo-Anosov Diffeomorphisms

Dominic Veconi

Pseudo-Anosov Maps

Smooth Ergodic Theory

Young Towers

# Behavior at Singularities

► The orthonormal basis are tangent vectors of the stable and unstable leaves. Along different prongs of the singularities, matrix form of *df* approaches different rotations of (<sup>λ</sup><sub>0</sub> <sup>0</sup><sub>λ<sup>-1</sup></sub>). In particular, *f* is not differentiable at the singularities.



Thermodynamics of Pseudo-Anosov Diffeomorphisms

Dominic Veconi

Pseudo-Anosov Maps

Smooth Ergodic Theory

Young Towers

# Simplifying assumptions

Since f is a homeomorphism, f permutes the singularities, so we may assume singularities are fixed points. Furthermore, we may assume that near singularities, the open sectors between the stable prongs are invariant under f (see the colored sections below).

Thermodynamics of Pseudo-Anosov Diffeomorphisms

Dominic Veconi

Pseudo-Anosov Maps

Smooth Ergodic Theory

Young Towers

## Slow-down procedure

Each open sector is homeomorphic to the right half-plane, where in coordinates we have s<sub>1</sub> = const are the stable leaves, and s<sub>2</sub> = const are unstable leaves.



In these coordinates, the map has the form

$$f(s_1, s_2) = (\lambda s_1, \lambda^{-1} s_2) \tag{1}$$

which is the time-1 map of the flow given by

$$\dot{s}_1 = s_1 \log \lambda, \quad \dot{s}_2 = -s_2 \log \lambda.$$

Thermodynamics of Pseudo-Anosov Diffeomorphisms

Dominic Veconi

Pseudo-Anosov Maps

Smooth Ergodic Theory

Young Towers

Proof of Main Result

◆□▶ ◆□▶ ◆三▶ ◆三▶ ○○○

#### Slow-down procedure

- For each singularity x<sub>i</sub>, choose coordinate ball of radius r<sub>i</sub> > 0 in which f(s<sub>1</sub>, s<sub>2</sub>) = (λs<sub>1</sub>, λ<sup>-1</sup>s<sub>2</sub>) in each sector. Assume r<sub>i</sub> = r<sub>j</sub> whenever x<sub>i</sub> and x<sub>j</sub> have the same number of prongs.
- ▶ Let  $0 < \tilde{r_i} < r_i$ . Suppose  $x_i$  has p prongs. Define a "slow-down" function  $\Psi_p : [0, \infty) \to \mathbb{R}$  so that:

1. 
$$\Psi_{p}(u) = C_{p}u^{(p-2)/p}$$
 for  $u \leq \tilde{r}_{i}^{2}$ , where  $C_{p} = (p/2)^{(2p-4)/p}$ ;  
2.  $\Psi_{p}$  is  $C^{\infty}$  except at 0;  
3.  $\dot{\Psi}_{p}(u) \geq 0$  for  $u > 0$ ;  
4.  $\Psi_{p}(u) = 1$  for  $u \geq r_{i}^{2}$ .

Let G<sub>p</sub> be the time-1 map of the flow given by the vector field defined by

$$\begin{cases} \dot{s}_1 = (\log \lambda) s_1 \Psi_p \left( s_1^2 + s_2^2 \right), \\ \dot{s}_2 = -(\log \lambda) s_2 \Psi_p \left( s_1^2 + s_2^2 \right). \end{cases}$$
(2)

Thermodynamics of Pseudo-Anosov Diffeomorphisms

Dominic Veconi

Pseudo-Anosov Maps

Smooth Ergodic Theory

Young Towers

Proof of Main Result

(ロ)、

## Slow-down procedure

▶ In coordinates, for  $|(s_1, s_2)| \ge r_i$ , we have  $G_p(s_1, s_2) = f(s_1, s_2)$ , so we define  $g : M \to M$  in coordinates by

$$g(x) = \begin{cases} G_p(s_1, s_2) & \text{if } x = (s_1, s_2) \text{ is near a singularity,} \\ f(x) & \text{otherwise.} \end{cases}$$

- Compare to the Katok map G : T<sup>2</sup> → T<sup>2</sup>, which is a toral automorphism that has similarly been slowed down at the origin.
  - After slow-down, the Katok map is conjugated with a homeomorphism to make the map Lebesgue-preserving ("blows up" trajectories near the origin).
- For smooth pseudo-Anosov maps, we instead show g preserves the measure Ψ<sub>p</sub>(s<sub>1</sub><sup>2</sup> + s<sub>2</sub><sup>2</sup>)<sup>-1</sup>ds<sub>1</sub> ∧ ds<sub>2</sub>.

Thermodynamics of Pseudo-Anosov Diffeomorphisms

Dominic Veconi

Pseudo-Anosov Maps

Smooth Ergodic Theory

Young Towers

# Pseudo-Anosov Diffeomorphisms

#### Theorem (Gerber and Katok, 1982)

- ► The map g is a C<sup>∞</sup> nonuniformly hyperbolic diffeomorphism of M.
- g is topologically conjugate to the pseudo-Anosov map f via a homeomorphism that is isotopic to the identity.
- This conjugacy is a homeomorphism only, and cannot be made C<sup>1</sup>.
- In every neighborhood of the singularities, g is real analytic, and is Bernoulli with respect to an invariant measure given by a smooth positive density.

Thermodynamics of Pseudo-Anosov Diffeomorphisms

Dominic Veconi

Pseudo-Anosov Maps

Smooth Ergodic Theory

Young Towers

# Equilibrium states and geometric potentials

• Let  $\varphi : M \to \mathbb{R}$  be continuous. A probability measure  $\mu_{\varphi}$  is an **equilibrium measure** for  $\varphi$  if

$$P_g(\varphi) = h_{\mu_{\varphi}}(g) + \int_M \varphi \, d\mu_{\varphi},$$

where  $h_{\mu_{\varphi}}(g)$  is the metric entropy of g and  $P_g(\varphi)$  is the topological pressure of  $\varphi$ :

$$P_g(\varphi) = \sup_{\mu \in \mathcal{M}(g)} \left\{ h_\mu(g) + \int_M \varphi \, d\mu \right\}$$

 We consider equilibrium states of the geometric t-potential

$$\varphi_t(x) = -t \log \left| dg \right|_{E^u(x)} \right|.$$

We denote  $\mu_t := \mu_{\varphi_t}$ .

• Observe that  $\mu_0$  is a measure of maximal entropy.

Thermodynamics of Pseudo-Anosov Diffeomorphisms

Dominic Veconi

Pseudo-Anosov Maps

Smooth Ergodic Theory

Young Towers

## Decay of correlations and CLT

*f* has exponential decay of correlations with respect to a measure µ and a class of functions H on M if there exists κ ∈ (0, 1) s.t. for any h<sub>1</sub>, h<sub>2</sub> ∈ H,

$$\left|\int (h_1 \circ f^n) h_2 d\mu - \int h_1 d\mu \int h_2 d\mu\right| \leq C \kappa^n$$

for some  $C = C(h_1, h_2) > 0$ .

*f* satisfies the Central Limit Theorem (CLT) if for any *h* ∈ H s.t. *h* ≠ *h*′ ∘ *f* − *h*′, *h*′ ∈ H, there is σ > 0 s.t.

$$\lim_{n\to\infty} \mu \left\{ \sqrt{n} \left( \frac{1}{n} S_n(h) - \mathbb{E}(h) \right) < t \right\}$$
$$= \frac{1}{\sigma \sqrt{2\pi}} \int_{-\infty}^t e^{-\tau^2/2\sigma^2} d\tau$$

where  $S_n(h) = \sum_{i=0}^{n-1} h(f^i(x))$  and  $\mathbb{E}(h) = \int_M h \, d\mu$ .

Thermodynamics of Pseudo-Anosov Diffeomorphisms

Dominic Veconi

Pseudo-Anosov Maps

Smooth Ergodic Theory

Young Towers

# Main Result

#### Theorem (V. 2020)

Let  $g : M \to M$  be a pseudo-Anosov diffeomorphism of a compact orientable manifold M (as in the preceding construction).

- 1. For any  $t_0 < 0$ , we may choose radii  $r_i > 0$  in the construction of g s.t. for  $t \in (t_0, 1)$ , there is a unique equilibrium measure  $\mu_t$  for the geometric potential  $\varphi_t$ . Further:
  - μ<sub>t</sub> satisfies CLT with respect to a class of functions containing all Hölder functions;
  - μ<sub>t</sub> has exponential decay of correlations with respect to this class of functions, and is hence mixing;
  - the map is Bernoulli with respect to  $\mu_t$ ;
  - the pressure function  $t \mapsto P_g(\varphi_t)$  is real-analytic on  $(t_0, 1)$ .

Thermodynamics of Pseudo-Anosov Diffeomorphisms

Dominic Veconi

Pseudo-Anosov Maps

Smooth Ergodic Theory

Young Towers

# Main Result (cont)

- 2. For t = 1, there are two classes of equilibrium measures associated to  $\varphi_1$ :
  - convex combinations of the Dirac measures δ<sub>xi</sub> centered at the singularities, and
  - a unique invariant SRB measure.
- 3. For t > 1, all equilibrium measures for  $\varphi_t$  are convex combinations of the measures  $\delta_{x_i}$ .

This result closely mirrors a similar result (Pesin, Senti, and Zhang, 2017) about the Katok map  $G : \mathbb{T}^2 \to \mathbb{T}^2$ .

 Replace "convex combinations of δ<sub>xi</sub>" with "the Dirac measure at the origin". Thermodynamics of Pseudo-Anosov Diffeomorphisms

Dominic Veconi

Pseudo-Anosov Maps

Smooth Ergodic Theory

Young Towers

# Young diffeomorphisms (general idea)

- The proof of the main result relies on the technology of Young towers.
- Given g : M → M and Λ ⊂ M, let τ : Λ → N be an inducing time (often first-return time) and let G = g<sup>τ</sup> : Λ → Λ be the induced map, defined by G(x) = g<sup>τ(x)</sup>(x).
- The map g : M → M is a Young diffeomorphism with base Λ ⊂ M if Λ has hyperbolic product structure, and G satisfies certain "nice" properties, including:
  - Stable (resp. unstable) leaves are invariant under G (resp. G<sup>-1</sup>);
  - G (resp. G<sup>-1</sup>) contracts points in the same stable (resp. unstable) leaf as n→∞ (resp. n→-∞);
  - τ is integrable on some unstable leaf;
  - Distortion estimates are bounded (more on this later).

Thermodynamics of Pseudo-Anosov Diffeomorphisms

Dominic Veconi

Pseudo-Anosov Maps

Smooth Ergodic Theory

Young Towers

# Thermodynamics of Young's diffeomorphisms

Let g : M → M be a C<sup>1+ε</sup> Young diffeomorphism of a compact Riemannian manifold M with base Λ ⊂ M and first return time τ : Λ → N. Under certain arithmetic and combinatorial conditions:

#### Theorem (Pesin, Senti, Zhang 2016)

- ►  $\exists$  an equilibrium measure  $\mu_1$  for the potential  $\varphi_1$ , which is the unique SRB measure;
- ► ∃  $t_0 < 0$  s.t. for  $t \in (t_0, 1)$ , there is a unique equilibrium measure  $\mu_t$  for  $\varphi_t$  on  $Y := \{g^k(x) : x \in \Lambda, 0 \le k \le \tau(x) - 1\};$
- For t ∈ (t<sub>0</sub>, 1), the measure µ<sub>t</sub> has exponential decay of correlations and the CLT with respect to a class of functions ℋ containing all Hölder functions on M.

#### Theorem (Shahidi, Zelerowicz 2018)

If  $g: M \to M$  is mixing, then  $(M, g, \mu_t)$  is Bernoulli.

Thermodynamics of Pseudo-Anosov Diffeomorphisms

Dominic Veconi

Pseudo-Anosov Maps

Smooth Ergodic Theory

Young Towers

# Constructing Tower

- ▶ Let  $\mathcal{P}$  be a Markov partition for g, and let  $P \in \mathcal{P}$  be a rectangle that does not contain any singularity.
- Let  $\tau(x)$  be first return time of x to P.
- For x ∈ P, let γ<sup>s</sup>(x) and γ<sup>u</sup>(x) be the connected component of the intersection of the stable and unstable leaves with P.
- For x with τ(x) < ∞, let U<sup>u</sup>(x) ⊆ γ<sup>u</sup>(x) be an open interval containing x, and

$$A^u(x) = \{y \in U^u(x) : y \in \partial P \text{ or } \tau(y) = \infty\}.$$

Assume  $U^{u}(x)$  is small enough s.t.  $\tau|_{U^{u}(x)\setminus A^{u}(x)} \equiv \text{const } \forall x \in P \text{ w} / \tau(x) < \infty.$  $\blacktriangleright$  Define the "stable strips":

$$\Lambda^{s}(x) = \bigcup_{y \in U^{u}(x) \setminus A^{u}(x)} \gamma^{s}(y).$$

Thermodynamics of Pseudo-Anosov Diffeomorphisms

Dominic Veconi

Pseudo-Anosov Maps

Smooth Ergodic Theory

Young Towers

Proof of Main Result

◆□▶ ◆□▶ ◆臣▶ ◆臣▶ 臣 のへで

# Constructing Tower

► We get countable collection  $\{\Lambda_i^s\}_{i\geq 1} \le \tau/\tau \mid_{\Lambda_i^s} \equiv \tau_i \in \mathbb{N}$ . Define  $\Lambda = \bigcup_{i\geq 1} \Lambda_i^s$ .



Thermodynamics of Pseudo-Anosov Diffeomorphisms

Dominic Veconi

Pseudo-Anosov Maps

Smooth Ergodic Theory

Young Towers

Proof of Main Result

#### Theorem (V. 2020)

The smooth pseudo-Anosov diffeomorphism  $g : M \to M$  is a Young's diffeomorphism with tower base  $\Lambda$ .

# **Bounded Distortion**

Most properties of Young diffeomorphisms are easy to verify, and follow from corresponding properties of pseudo-Anosov diffeomorphisms. The one tricky property is *bounded distortion*:

#### Lemma

There exist c > 0 and  $\kappa \in (0, 1)$  such that for all  $n \ge 0$ ,  $x \in \Lambda$  and  $y \in \gamma^{s}(x)$ , we have

$$\left|\log\frac{\left|dG|_{E^u(G^n(x))}\right|}{\left|dG|_{E^u(G^n(y))}\right|}\right| \leq c\kappa^n.$$

This bound is easy to show outside of slow-down neighborhoods. Inside the slow-down, there is a bound on how far apart log |dg<sub>E<sup>u</sup>(g<sup>n</sup>(x))</sub>| and log |dg<sub>E<sup>u</sup>(g<sup>n</sup>(y))</sub>| can be. (This is why we assume stable sectors are locally invariant.)

Thermodynamics of Pseudo-Anosov Diffeomorphisms

Dominic Veconi

Pseudo-Anosov Maps

Smooth Ergodic Theory

Young Towers

## Equilibrium state existence

► Using previous results, this gives us a unique equilibrium measure µ<sub>t</sub> for t < 1 on the set</p>

$$Y := \left\{ g^k(x) : x \in \Lambda, 0 \le k \le \tau(x) - 1 \right\}$$

- If P̂ is another element of the Markov partition for (M, g), same argument gives us unique equilibrium measure µ̂t for t < 1 and corresponding set Ŷ.</p>
- Assuming (M, g) is topologically transitive, since µ<sub>t</sub>(U) > 0 and µ̂<sub>t</sub>(Û) > 0 for every open U ⊃ P, Û ⊃ P̂, and g<sup>k</sup>(U) ∩ Û ≠ Ø for some k ≥ 1, it follows from uniqueness that µ<sub>t</sub> = µ̂<sub>t</sub>.

Thermodynamics of Pseudo-Anosov Diffeomorphisms

Dominic Veconi

Pseudo-Anosov Maps

Smooth Ergodic Theory

Young Towers

#### The t = 1 case

For t = 1, we get at least one equilibrium measure, μ<sub>1</sub>, which is an SRB measure. By the Pesin entropy formula,

$$P_{g}(\varphi_{1}) = h_{\mu_{1}}(g) - \int_{M} \log |dg|_{E^{u}(x)}| d\mu_{1}(x) = 0.$$

- If ν is any other equilibrium measure for φ<sub>1</sub>, it also satisfies the entropy formula. So if ν has positive Lyapunov exponents, ν is an SRB measure. By uniqueness of SRB measures, ν = μ<sub>1</sub>.
- But if ν has no positive Lyapunov exponents, then log |dg|<sub>E<sup>u</sup>(x)</sub>| = 0 ν-a.e., so ν is supported on the (finite) set of singularities. So ν = ∑ α<sub>i</sub>δ<sub>xi</sub> is a convex combination of the Dirac measures on the singularities. (If t > 1, can be similarly showed that all equilibrium states are combinations of Dirac measures.)

Thermodynamics of Pseudo-Anosov Diffeomorphisms

Dominic Veconi

Pseudo-Anosov Maps

Smooth Ergodic Theory

Young Towers

# Further results

# Theorem (Pesin, Senti, Zhang 2018)

The Katok map  $G : \mathbb{T}^2 \to \mathbb{T}^2$  has polynomial decay of correlations with respect to its unique SRB measure:

$$\left|\int (h_1 \circ f^n) h_2 d\mu - \int h_1 d\mu \int h_2 d\mu\right| \leq C n^{-\kappa}$$

#### Theorem (Wang 2020)

The Katok map G has a unique equilibrium state for  $\varphi_t$ , for every t < 1.

**Question:** Is this also true for pseudo-Anosov diffeomorphisms?

Both of these results assume the exponent α > 0 in the slowing down of the Katok map is < 1/2. Our exponent is (p − 2)/p > 1/2 when p ≥ 5.

Thermodynamics of Pseudo-Anosov Diffeomorphisms

Dominic Veconi

Pseudo-Anosov Maps

Smooth Ergodic Theory

Young Towers

#### References

- A. Fathi, F. Laudenbach, and V. Poénaru, Travaux de Thurston sur les surfaces, in Séminaire Orsay, Astérisque 66-67, 1979.
- M. Gerber, A. Katok. Smooth models of pseudo-Anosov maps. Ann. scient. Éc. Norm. Sup. 4(15):173-204, 1982.
- E. Lanneau. Tell me a pseudo-Anosov. EMS Newsletter, Dec. 2017, 12-16.
- F. Ledrappier and L. S. Young. The metric entropy of diffeomorphisms, Bull. of the Amer. Math. Soc. (N.S.), 11(2):343-346, 1984.
- Y. Pesin, S. Senti, and K. Zhang. Thermodynamics of the Katok map (revised version). *Ergod. Theory Dyn. Syst.*, **39**(3):764-794, 2019
- Y. Pesin, S. Senti, and K. Zhang. Thermodynamics of towers of hyperbolic type. Trans. Amer. Math. Soc. 368(12):8519-8552. 2016.
- F. Rodriguez-Hertz, M. A. Rodriguez-Hertz, A. Tahzibi, and R. Ures, Uniqueness of SRB measures for transitive diffeomorphisms on surfaces, *Commun. Math. Phys.*, **306**(1):35-49, 2011.
- F. Shahidi, A. Zelerowicz. Thermodynamics via inducing, J. Stat. Phys, 175(2):351-383, 2019.
- D. Veconi. Thermodynamics of Smooth Models of Pseudo-Anosov Homeomorphisms. Preprint.
- T. Wang. Unique equilibrium states, large deviations and Lyapunov spectra for the Katok map. *Ergod. Theory Dyn. Syst.*, 2020.

Thermodynamics of Pseudo-Anosov Diffeomorphisms

Dominic Veconi

Pseudo-Anosov Maps

Smooth Ergodic Theory

Young Towers