

Thermodynamics of Pseudo-Anosov Diffeomorphisms

University of Maryland Dynamical Systems Seminar

Dominic Veconi

Penn State University

October 15 2020

Why study these?

- ▶ *Smooth ergodic theory*—i.e. the ergodic theory of smooth (often hyperbolic) maps.
- ▶ A diffeomorphism $f : M \rightarrow M$ of a Riemannian manifold is **Anosov** if at every $x \in M$, the tangent space splits into two invariant subspaces $T_x M = E^u(x) \oplus E^s(x)$:

$$\begin{aligned} |df_x^n v| &\leq \lambda^{-n} |v| && \text{for every } v \in E^s(x); && \text{and} \\ |df_x^{-n} v| &\leq \lambda^{-n} |v| && \text{for every } v \in E^u(x). \end{aligned}$$

- ▶ In low dimensions, these are hyperbolic toral automorphisms, $A : \mathbb{T}^2 \rightarrow \mathbb{T}^2$, $A \in \text{SL}(2, \mathbb{Z})$, and their perturbations.
- ▶ If we can't have Anosov maps on other surfaces, what's the next best thing?

Pseudo-Anosov Homeomorphisms

- ▶ A homeomorphism $f : M \rightarrow M$ of a 2-manifold M is *pseudo-Anosov* if there are two f -invariant foliations with singularities, \mathcal{F}^s and \mathcal{F}^u , for which:
 1. the foliations share the same singularities, and the same number of prongs;
 2. the foliations intersect transversally away from the singularities;
 3. there is a $\lambda > 1$ such that for x, y in the same \mathcal{F}^s -leaf, $\rho^s(f(x), f(y)) = \lambda^{-1}\rho^s(x, y)$, and for x, y in the same \mathcal{F}^u -leaf, $\rho^u(f(x), f(y)) = \lambda\rho^u(x, y)$;

where ρ^s and ρ^u are the distances in the \mathcal{F}^s and \mathcal{F}^u foliations with respect to a Riemannian metric on M that has a density vanishing at the singularities.

Nielsen-Thurston Classification

Theorem (Nielsen, Thurston)


Any homeomorphism on a compact topological manifold M is isotopic to a map f that is one of the following:

- ▶ f is **periodic**: there is a positive integer m with $f^m = \text{Id}$;
 - ▶ EG. $f = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} : \mathbb{T}^2 \rightarrow \mathbb{T}^2$;
- ▶ f is **reducible**: there is a closed curve on M that is f -invariant (these are also known as Dehn twists);
 - ▶ EG. $f = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} : \mathbb{T}^2 \rightarrow \mathbb{T}^2$;
- ▶ f is **pseudo-Anosov**;
 - ▶ EG. $f = \begin{pmatrix} 2 & 1 \\ 1 & 1 \end{pmatrix} : \mathbb{T}^2 \rightarrow \mathbb{T}^2$.

The pseudo-Anosov maps form an open set in the homeomorphisms of M , and exhibit the most interesting dynamical properties.


Examples from Anosov maps

- ▶ An Anosov diffeomorphism is a pseudo-Anosov homeomorphism with no singularities.
- ▶ Linear Anosov maps on \mathbb{T}^2 lift to pseudo-Anosov maps on higher-genus surfaces via branched coverings (may be necessary to lift powers of Anosov maps).




Example from Dehn twists

- ▶ Orientable genus-2 surface S_2 can be horizontally split into two cylinders C_1 and C_2 , each of which admits a Dehn twist T_1 and T_2 resp. Note $dT_1 = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ on C_1 , and $dT_2 = \begin{pmatrix} 1 & 2 \\ 0 & 1 \end{pmatrix}$ on C_2 , away from the identified vertex (singularity).




Example from Dehn twists

- ▶ The Dehn twists T_1 and T_2 can be composed to form a horizontal “multi-twist” $T := T_1^2 \circ T_2$, whose differential away from the vertex is $\begin{pmatrix} 1 & 2 \\ 0 & 1 \end{pmatrix}$:




Example from Dehn twists

- ▶ A similar vertical multi-twist S can be defined on the cylinder made from the red and blue squares, and the cylinder made from the green square. The corresponding differential is $\begin{pmatrix} 1 & 0 \\ 2 & 1 \end{pmatrix}$.



Example from Dehn twists

- ▶ $T \circ S$ has the constant differential $\begin{pmatrix} 1 & 2 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 2 & 1 \end{pmatrix} = \begin{pmatrix} 5 & 2 \\ 2 & 1 \end{pmatrix}$.
- ▶ Eigenvalues are $3 - 2\sqrt{2}$ and $3 + 2\sqrt{2}$, with resp. eigenvectors $\begin{pmatrix} 1-\sqrt{2} \\ 1 \end{pmatrix}$ and $\begin{pmatrix} 1+\sqrt{2} \\ 1 \end{pmatrix}$.
- ▶ This is a pseudo-Anosov map whose stable/unstable foliations are parallel to the eigendirections for $3 - 2\sqrt{2}$ and $3 + 2\sqrt{2}$, resp. The vertex is a 6-pronged singularity (unstable prongs illustrated:)



- ▶ In fact, this map is the lift of the linear Anosov map on \mathbb{T}^2 induced by $\begin{pmatrix} 5 & 2 \\ 2 & 1 \end{pmatrix}$.

Suppose $f : M \rightarrow M$ is a pseudo-Anosov homeomorphism with expansion factor $\lambda > 1$.


- ▶ There is a Riemannian metric on M inducing a volume ν under which f is invariant, given locally by the product of ν^s and ν^u .
- ▶ If U is a neighborhood of a singularity $x_j \in M$ and $\varphi : U \rightarrow \mathbb{R}^2$ is a coordinate chart, ν has a density with respect to $\varphi_*^{-1}(\text{Leb})$ vanishing at x_j .
- ▶ If $x \in M$ is not a singularity, then f is smooth at x and there are orthonormal bases of $T_x M$ and $T_{f(x)} M$ with respect to which df_x has the matrix form $\begin{pmatrix} \lambda & 0 \\ 0 & \lambda^{-1} \end{pmatrix}$.

For proofs, see Fathi, Laudenbach, Poénaru, et al.

Thurston's Work on Surfaces.


Behavior at Singularities

- ▶ The orthonormal basis are tangent vectors of the **stable** and **unstable** leaves. Along different prongs of the singularities, matrix form of df approaches different rotations of $\begin{pmatrix} \lambda & 0 \\ 0 & \lambda^{-1} \end{pmatrix}$. In particular, f is not differentiable at the singularities.



Slow-down procedure

- ▶ Each open sector is homeomorphic to the right half-plane, where in coordinates we have $s_1 = \text{const}$ are the stable leaves, and $s_2 = \text{const}$ are unstable leaves.



- ▶ In these coordinates, the map has the form

$$f(s_1, s_2) = (\lambda s_1, \lambda^{-1} s_2) \quad (1)$$

which is the time-1 map of the flow given by

$$\dot{s}_1 = s_1 \log \lambda, \quad \dot{s}_2 = -s_2 \log \lambda.$$

Slow-down procedure

- ▶ For each singularity x_i , choose coordinate ball of radius $r_i > 0$ in which $f(s_1, s_2) = (\lambda s_1, \lambda^{-1} s_2)$ in each sector. Assume $r_i = r_j$ whenever x_i and x_j have the same number of prongs.
- ▶ Let $0 < \tilde{r}_i < r_i$. Suppose x_i has p prongs. Define a “slow-down” function $\Psi_p : [0, \infty) \rightarrow \mathbb{R}$ so that:
 1. $\Psi_p(u) = C_p u^{(p-2)/p}$ for $u \leq \tilde{r}_i^2$, where $C_p = (p/2)^{(2p-4)/p}$;
 2. Ψ_p is C^∞ except at 0;
 3. $\dot{\Psi}_p(u) \geq 0$ for $u > 0$;
 4. $\Psi_p(u) = 1$ for $u \geq r_i^2$.
- ▶ Let G_p be the time-1 map of the flow given by the vector field defined by

$$\begin{cases} \dot{s}_1 = (\log \lambda) s_1 \Psi_p(s_1^2 + s_2^2), \\ \dot{s}_2 = -(\log \lambda) s_2 \Psi_p(s_1^2 + s_2^2). \end{cases} \quad (2)$$

- ▶ Same trajectories as $\dot{s}_1 = (\log \lambda) s_1$, $\dot{s}_2 = -(\log \lambda) s_2$, but slower.

Slow-down procedure

- ▶ In coordinates, for $|(s_1, s_2)| \geq r_i$, we have $G_p(s_1, s_2) = f(s_1, s_2)$, so we define $g : M \rightarrow M$ in coordinates by

$$g(x) = \begin{cases} G_p(s_1, s_2) & \text{if } x = (s_1, s_2) \text{ is near a singularity,} \\ f(x) & \text{otherwise.} \end{cases}$$

- ▶ Compare to the *Katok map* $G : \mathbb{T}^2 \rightarrow \mathbb{T}^2$, which is a toral automorphism that has similarly been slowed down at the origin.
 - ▶ After slow-down, the Katok map is conjugated with a homeomorphism to make the map Lebesgue-preserving (“blows up” trajectories near the origin).
- ▶ For smooth pseudo-Anosov maps, we instead show g preserves the measure $\Psi_p(s_1^2 + s_2^2)^{-1} ds_1 \wedge ds_2$.

Theorem (Gerber and Katok, 1982)

- ▶ *The map g is a C^∞ nonuniformly hyperbolic diffeomorphism of M .*
- ▶ *g is topologically conjugate to the pseudo-Anosov map f via a homeomorphism that is isotopic to the identity.*
- ▶ *This conjugacy is a homeomorphism only, and cannot be made C^1 .*
- ▶ *In every neighborhood of the singularities, g is real analytic. Furthermore, g is Bernoulli with respect to an invariant measure given by a smooth positive density.*

Physical measures

- ▶ If we're studying ergodic theory, what measures are we interested in?
- ▶ Primarily we want to study the *physical measures*:

$$\mu \left\{ x : \frac{1}{n} \sum_{k=0}^{n-1} (\varphi \circ f^k)(x) \rightarrow \int \varphi d\mu \quad \forall \varphi \in C^0 \right\} > 0$$

- ▶ For hyperbolic systems $f : M \rightarrow M$, one important physical measures are the **Sinai-Ruelle-Bowen (SRB)** measures, which are f -invariant probability measures μ for which:
 - ▶ μ has positive Lyapunov exponents almost everywhere, and
 - ▶ μ has absolutely continuous conditional measures on unstable manifolds (w.r.t. Riemannian leaf volume).
- ▶ For Anosov and pseudo-Anosov maps that preserve area, SRB measures are simply this area.

Equilibrium states and geometric potentials

- ▶ Let $\varphi : M \rightarrow \mathbb{R}$ be continuous. A probability measure μ_φ is an **equilibrium measure** for φ if

$$P_g(\varphi) = h_{\mu_\varphi}(g) + \int_M \varphi d\mu_\varphi,$$

where $h_{\mu_\varphi}(g)$ is the metric entropy of g and $P_g(\varphi)$ is the topological pressure of φ :

$$P_g(\varphi) = \sup_{\mu \in \mathcal{M}(g)} \left\{ h_\mu(g) + \int_M \varphi d\mu \right\}$$

- ▶ We consider equilibrium states of the *geometric t -potential*

$$\varphi_t(x) = -t \log |dg|_{E^u(x)}|.$$

We denote $\mu_t := \mu_{\varphi_t}$.

- ▶ Observe that μ_0 is a measure of maximal entropy.

Ergodic theory of Anosov maps

- ▶ The special case $t = 1$ gives the *geometric potential*
 $\varphi_1(x) = -\log \det |dg|_{E^u(x)}|.$

Theorem

If $f : M \rightarrow M$ is an Anosov (or more generally Axiom A) diffeomorphism, there exists a unique SRB measure for f .

Proof.

From Bowen's notes:

- ▶ Any Hölder continuous potential $\varphi : M \rightarrow \mathbb{R}$ has a unique equilibrium state μ_φ for φ .
- ▶ The geometric potential $\varphi_1(x)$ is Hölder, so $\mu_{\varphi_1} = \mu_1$ has a unique equilibrium state.
- ▶ This equilibrium state satisfies all of the requisite properties of SRB measures.



The pseudo-Anosov case

- ▶ Bowen's proof of this result relies on two key points:
 - ▶ The map admits a finite Markov partition.
 - ▶ Geometric potential φ_1 is Hölder.

Proposition (Fathi, Shub '79; Katok, Gerber '82)

The pseudo-Anosov homeomorphism $f : M \rightarrow M$ admits a finite Markov partition, with respect to which f is Bernoulli.

- ▶ For “linear” pseudo-Anosov homeomorphisms, this argument applies basically verbatim.
- ▶ For the slowed-down Katok-Gerber diffeomorphisms, φ_1 is not Hölder (in particular its induced map on the finite symbolic space is not Hölder).

Decay of correlations and CLT

- ▶ f has **exponential decay of correlations** with respect to a measure μ and a class of functions \mathcal{H} on M if there exists $\kappa \in (0, 1)$ s.t. for any $h_1, h_2 \in \mathcal{H}$,

$$\left| \int (h_1 \circ f^n) h_2 d\mu - \int h_1 d\mu \int h_2 d\mu \right| \leq C \kappa^n$$

for some $C = C(h_1, h_2) > 0$.

- ▶ f satisfies the **Central Limit Theorem** (CLT) if for any $h \in \mathcal{H}$ s.t. $h \neq h' \circ f - h'$, $h' \in \mathcal{H}$, there is $\sigma > 0$ s.t.

$$\begin{aligned} \lim_{n \rightarrow \infty} \mu \left\{ \sqrt{n} \left(\frac{1}{n} S_n(h) - \mathbb{E}(h) \right) < t \right\} \\ = \frac{1}{\sigma \sqrt{2\pi}} \int_{-\infty}^t e^{-\tau^2/2\sigma^2} d\tau \end{aligned}$$

where $S_n(h) = \sum_{i=0}^{n-1} h(f^i(x))$ and $\mathbb{E}(h) = \int_M h d\mu$.

Theorem (V. 2020)

Let $g : M \rightarrow M$ be a pseudo-Anosov diffeomorphism of a compact orientable manifold M (as in the preceding construction).

1. For any $t_0 < 0$, we may choose radii $r_i > 0$ in the construction of g s.t. for $t \in (t_0, 1)$, there is a unique equilibrium measure μ_t for the geometric potential φ_t .
Further:

- ▶ μ_t satisfies CLT with respect to a class of functions containing all Hölder functions;
- ▶ μ_t has exponential decay of correlations with respect to this class of functions, and is hence mixing;
- ▶ the map is Bernoulli with respect to μ_t ;
- ▶ the pressure function $t \mapsto P_g(\varphi_t)$ is real-analytic on $(t_0, 1)$.

Main Result (cont)

2. For $t = 1$, there are two classes of equilibrium measures associated to φ_1 :
 - ▶ convex combinations of the Dirac measures δ_{x_i} centered at the singularities, and
 - ▶ a unique invariant SRB measure.
3. For $t > 1$, all equilibrium measures for φ_t are convex combinations of the measures δ_{x_i} .

This result closely mirrors a similar result (Pesin, Senti, and Zhang, 2017) about the Katok map $G : \mathbb{T}^2 \rightarrow \mathbb{T}^2$.

- ▶ Replace “convex combinations of δ_{x_i} ” with “the Dirac measure at the origin”.

Further results

Theorem (Pesin, Senti, Shahidi 2020)

The Katok map $G : \mathbb{T}^2 \rightarrow \mathbb{T}^2$ has polynomial decay of correlations with respect to Lebesgue area:

$$\left| \int (h_1 \circ f^n) h_2 d\mu - \int h_1 d\mu \int h_2 d\mu \right| \leq Cn^{-\kappa}$$

Theorem (Wang 2020)

The Katok map G has a unique equilibrium state for φ_t , for every $t < 1$.

Question: Is this also true for pseudo-Anosov diffeomorphisms?

- ▶ Both of these results assume the exponent $\alpha > 0$ in the slowing down of the Katok map is $< 1/2$. Our exponent is $(p - 2)/p > 1/2$ when $p \geq 5$.

Young diffeomorphisms (general idea)

- ▶ The proof of the main result relies on the technology of *Young towers*.
- ▶ Given $g : M \rightarrow M$ and $\Lambda \subset M$, let $\tau : \Lambda \rightarrow \mathbb{N}$ be an *inducing time* (often first-return time) and let $G = g^\tau : \Lambda \rightarrow \Lambda$ be the *induced map*, defined by $G(x) = g^{\tau(x)}(x)$.
- ▶ The map $g : M \rightarrow M$ is a *Young diffeomorphism* with base $\Lambda \subset M$ if Λ has hyperbolic product structure, and G satisfies certain “nice” properties, including:
 - ▶ Stable (resp. unstable) leaves are invariant under G (resp. G^{-1});
 - ▶ G (resp. G^{-1}) contracts points in the same stable (resp. unstable) leaf as $n \rightarrow \infty$ (resp. $n \rightarrow -\infty$);
 - ▶ τ is integrable on some unstable leaf;
 - ▶ Distortion estimates are bounded (more on this later).

Thermodynamics of Young's diffeomorphisms

- ▶ Let $g : M \rightarrow M$ be a $C^{1+\varepsilon}$ Young diffeomorphism of a compact Riemannian manifold M with base $\Lambda \subset M$ and first return time $\tau : \Lambda \rightarrow \mathbb{N}$.

Theorem (Pesin, Senti, Zhang 2016)

- ▶ \exists an equilibrium measure μ_1 for the potential φ_1 , which is the unique SRB measure.
- ▶ Suppose for some $C > 0$, $0 < h < h_{\mu_1}(f)$, we have

$$S_n := \# \{ \Lambda_i^s : \tau_i = n \} \leq C e^{hn}.$$

$\exists t_0 < 0$ s.t. for $t \in (t_0, 1)$, there is a unique equilibrium measure μ_t for φ_t on

$$Y := \{ g^k(x) : x \in \Lambda, 0 \leq k \leq \tau(x) - 1 \}.$$

- ▶ For $t \in (t_0, 1)$, the measure μ_t has exponential decay of correlations and the CLT with respect to a class of functions \mathcal{H} containing all Hölder functions on M .

Constructing Tower

- ▶ Let \mathcal{P} be a Markov partition for g , and let $P \in \mathcal{P}$ be a rectangle that does not contain any singularity.
- ▶ Let $\tau(x)$ be first return time of x to P .
- ▶ For $x \in P$, let $\gamma^s(x)$ and $\gamma^u(x)$ be the connected component of the intersection of the stable and unstable leaves with P .
- ▶ For x with $\tau(x) < \infty$, let $U^u(x) \subseteq \gamma^u(x)$ be an open interval containing x , and

$$A^u(x) = \{y \in U^u(x) : y \in \partial P \text{ or } \tau(y) = \infty\}.$$

Assume $U^u(x)$ is small enough s.t.


$$\tau|_{U^u(x) \setminus A^u(x)} \equiv \text{const} \quad \forall x \in P \text{ w/ } \tau(x) < \infty.$$

- ▶ Define the “stable strips”:

$$\Lambda^s(x) = \bigcup_{y \in U^u(x) \setminus A^u(x)} \gamma^s(y).$$

Constructing Tower

- ▶ We get countable collection $\{\Lambda_i^s\}_{i \geq 1}$ w/ $\tau|_{\Lambda_i^s} \equiv \tau_i \in \mathbb{N}$.
Define $\Lambda = \bigcup_{i \geq 1} \Lambda_i^s$.



Theorem (V. 2020)

The smooth pseudo-Anosov diffeomorphism $g : M \rightarrow M$ is a Young's diffeomorphism with tower base Λ .

Bounded Distortion

- ▶ Most properties of Young diffeomorphisms are easy to verify, and follow from corresponding properties of pseudo-Anosov diffeomorphisms. The one tricky property is *bounded distortion*:

Lemma

There exist $c > 0$ and $\kappa \in (0, 1)$ such that for all $n \geq 0$, $x \in \Lambda$ and $y \in \gamma^s(x)$, we have

$$\left| \log \frac{|dG|_{E^u(G^n(x))}}{|dG|_{E^u(G^n(y))}} \right| \leq c\kappa^n.$$

- ▶ This bound is easy to show outside of slow-down neighborhoods. Inside the slow-down, there is a bound on how far apart $\log |dg_{E^u(g^n(x))}|$ and $\log |dg_{E^u(g^n(y))}|$ can be. (This is why we assume stable sectors are locally invariant.)

The $t = 1$ case

- ▶ For any equilibrium measure ν for $\varphi_1(x) = -\log \det |dg|_{E^u(x)}|$, by the Margulis-Ruelle Inequality,

$$P(\varphi_1) = h_\nu(g) + \int \varphi_1 d\nu \leq 0. \quad (3)$$

Theorem (Pesin, Senti, Zhang '16)

If $f : M \rightarrow M$ is a Young's diffeomorphism, there is a unique SRB measure for φ_1 .

- ▶ For the SRB measure/Riemannian volume, we have equality (by Pesin entropy formula).

The $t = 1$ case; other measures

Theorem (Ledrappier, Young '84)

Any measure with positive Lyapunov exponents satisfying Pesin entropy formula is an SRB measure.

Theorem (Rodriguez-Herz, Rodriguez-Herz, Tahzibi, Ures '11)

If $f : M \rightarrow M$ is a surface diffeomorphism admitting an SRB measure, then this SRB measure is unique.

- ▶ So if ν is any equilibrium measure with positive Lyapunov exponents, it is an SRB measure:

$$P(\varphi_1) = h_\nu(g) - \int \log \det |dg|_{E^u(x)}| d\nu(x) = 0$$

- ▶ But if ν has no positive Lyapunov exponents, then $\log |dg|_{E^u(x)}| = 0$ ν -a.e., so ν is supported on the (finite) set of singularities.
- ▶ So $\nu = \sum \alpha_i \delta_{x_i}$ is a convex combination of the Dirac measures on the singularities.

Equilibrium state existence for $t < 1$

- ▶ To use tower thermodynamics for $t < 1$, we need

$$S_n := \# \{ \Lambda_i^s : \tau_i = n \} \leq C e^{hn} \quad (4)$$

for some $C > 0$, $h < h_{\mu_1}(g)$.

Theorem (Fathi, Shub '79)

If $f : M \rightarrow M$ is a pseudo-Anosov homeomorphism with expansive constant $\lambda > 1$, and μ the Riemannian volume invariant under f , then $h_{\text{top}}(g) = h_{\text{top}}(f) = h_{\mu}(f) = \log \lambda$.

- ▶ Since f and g are topologically conjugate, $h_{\text{top}}(f) = h_{\text{top}}(g) = \log \lambda$.
- ▶ By studying symbolic systems induced by Markov partitions on f and g , (4) is satisfied for an $h < h_{\text{top}}(f) = h_{\text{top}}(g)$.

Equilibrium state existence for $t < 1$

- ▶ By taking slowdown neighborhoods to be sufficiently small, we can take $|h_{\mu_1}(g) - h_{\mu}(f)| < \varepsilon$ (μ_1 being the SRB for g).
- ▶ More explicitly, let \mathcal{U} be the union of the slowdown neighborhoods. If \mathcal{U} is sufficiently small, since $\det |dg|_{E^u(x)}| = \lambda$ and $\frac{d\mu}{d\mu_1} = 1$ outside of \mathcal{U} ,

$$\begin{aligned}h_{\mu}(f) - h_{\mu_1}(g) &= \int_M \left(\log \lambda \frac{d\mu}{d\mu_1} - \log \det |dg|_{E^u(x)} \right) d\mu_1 \\ &= \int_{\mathcal{U}} \left(\log \lambda \frac{d\mu}{d\mu_1} - \log \det |dg|_{E^u(x)} \right) d\mu_1 \\ &< \varepsilon.\end{aligned}$$

- ▶ It follows that $h_{\text{top}}(g) - h_{\mu_1}(g)$ may be made arbitrarily small. So $\exists h < h_{\mu_1}(g)$ satisfying $S_n \leq Ce^{hn}$.

Equilibrium state existence for $t < 1$

- ▶ Using previous results, this gives us a unique equilibrium measure μ_t for $t < 1$ on the set

$$Y := \left\{ g^k(x) : x \in \Lambda, 0 \leq k \leq \tau(x) - 1 \right\}$$

- ▶ If \hat{P} is another element of the Markov partition for (M, g) , same argument gives us unique equilibrium measure $\hat{\mu}_t$ for $t < 1$ and corresponding set \hat{Y} .
- ▶ Assuming (M, g) is topologically transitive, since $\mu_t(U) > 0$ and $\hat{\mu}_t(\hat{U}) > 0$ for every open $U \supset P$, $\hat{U} \supset \hat{P}$, and $g^k(U) \cap \hat{U} \neq \emptyset$ for some $k \geq 1$, it follows from uniqueness that $\mu_t = \hat{\mu}_t$.

The $t > 1$ case

- ▶ If $t > 1$ and ν an equilibrium state for φ_t , then again by Margulis-Ruelle,

$$h_\nu(g) \leq t \int \log \det |dg|_{E^u(x)}| d\nu(x)$$

w/ equality $\iff \int \log \det |dg|_{E^u(x)}| d\nu(x) = 0$.

- ▶ If δ is a convex combination of Dirac measures on singularities,

$$P(\varphi_t) = h_\nu(g) + \int \varphi_t d\nu \leq 0 = h_\delta(g) + \int \varphi_t d\delta$$

so we have equality, so $\log \det |dg|_{E^u(x)}| = 0$ ν -a.e., so ν is supported on the singularities.

References

- ▶ A. Fathi, F. Laudenbach, and V. Poénaru, Travaux de Thurston sur les surfaces, in Séminaire Orsay, *Astérisque* 66-67, 1979.
- ▶ M. Gerber, A. Katok. Smooth models of pseudo-Anosov maps. *Ann. scient. Éc. Norm. Sup.* **4**(15):173-204, 1982.
- ▶ E. Lanneau. Tell me a pseudo-Anosov. *EMS Newsletter*, Dec. 2017, 12-16.
- ▶ F. Ledrappier and L. S. Young. The metric entropy of diffeomorphisms, *Bull. of the Amer. Math. Soc. (N.S.)*, **11**(2):343-346, 1984.
- ▶ Y. Pesin, S. Senti, and K. Zhang. Thermodynamics of the Katok map (revised version). *Ergod. Theory Dyn. Syst.*, **39**(3):764-794, 2019
- ▶ Y. Pesin, S. Senti, and K. Zhang. Thermodynamics of towers of hyperbolic type. *Trans. Amer. Math. Soc.* **368**(12):8519-8552. 2016.
- ▶ F. Rodriguez-Hertz, M. A. Rodriguez-Hertz, A. Tahzibi, and R. Ures, Uniqueness of SRB measures for transitive diffeomorphisms on surfaces, *Commun. Math. Phys.*, **306**(1):35-49, 2011.
- ▶ F. Shahidi, A. Zelerowicz. Thermodynamics via inducing, *J. Stat. Phys.*, **175**(2):351-383, 2019.
- ▶ D. Veconi. Thermodynamics of Smooth Models of Pseudo-Anosov Homeomorphisms. Preprint.
- ▶ T. Wang. Unique equilibrium states, large deviations and Lyapunov spectra for the Katok map. *Ergod. Theory Dyn. Syst.*, 2020.